

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

Application number: 80902298.1

Date of filing: 25.11.80

Data of the international application taken as a basis:

International application number:
PCT/JP 80/00286

International publication number:
WO 81/01554 (11.06.81 81/14)

Int. Cl.²: **C 07 D 413/06, C 07 D 413/14,**
C 07 D 471/04, C 07 D 471/10,
C 07 D 471/20

Priority: 28.11.79 JP 154715/79

Date of publication of application: 02.12.81
Bulletin 81/48

Designated Contracting States: **DE FR GB NL**


Applicant: **Yoshitomi Pharmaceutical Industries, Ltd.,**
35 Hiranomachi 3-chome Higashi-ku, Osaka 541 (JP)

Inventor: **KAWAKITA, Takeshi, 7-8,**
Chuoma-chi 1-chome, Nakatsu-shi, Oita 871 (JP)
Inventor: **MURO, Tomio, 631-7, Oaza-kakize**
Nakatsu-shi, Oita 871 (JP)
Inventor: **SETOGUCHI, Michihide, 5-21,**
Chuomachi 2-chome, Nakatsu-shi, Oita 871 (JP)



Representative: **Heunemann, Dieter Dr., VOSSIUS**
VOSSIUS TAUCHNER HEUNEMANN RAUH P.O.
Box 86 07 67 Siebertstrasse 4, D-8000 Munich 86 (DE)

ISOXAZOLE DERIVATIVES.

Isoxazole derivatives represented by the following general formula:



[wherein Ar represents a phenyl or pyridyl group optionally substituted by halogen or lower alkoxy, R¹ represents a hydrogen atom, a lower alkyl group or a group of Ar, R² represents a hydrogen atom or, when R¹ and R² are taken together, they form a carbon-to-carbon bond, and Am represents an amine residue selected from the following:




wherein R³ represents a hydrogen atom or a lower alkyl-group, X¹ and X² each represent a hydrogen atom, a halogen atom or a trifluoromethyl group, and Y represents O or S] and the salts thereof. These are useful as psychotropic drugs and antimitotics.

Specification

Isoxazole derivatives










Technical field and Disclosure of the Invention

This invention relates to novel isoxazole derivatives having spontaneous locomotor suppressing activity, anti-apomorphine activity and like activity and useful as drugs such as psychotropic agent and antiemetic agent and represented by the general formula




and pharmaceutically acceptable acid addition salts thereof, and also to a process for preparing these compounds.

In the foregoing formula, Ar represents a phenyl group optionally containing a lower alkoxy group or a halogen atom as a substituent, or a pyridyl group, R¹ represents a hydrogen atom, a lower alkyl group or a group represented by Ar, R² represents a hydrogen atom or alternatively R¹ and R² are bound together and form a carbon-carbon bond, and Am represents an amino residue selected from the group consisting of the following residues:




and



wherein R^3 represents a hydrogen atom or a lower alkyl group, X^1 and X^2 each represent a hydrogen atom, a halogen atom or a trifluoromethyl group, and Y represents O or S.

The term "halogen" herein includes fluorine, chlorine, bromine, etc. The term "lower alkoxy" herein represents a methoxy, ethoxy, propoxy, butoxy, etc. The term "lower alkyl" herein represents methyl, ethyl, propyl, butyl, etc.

The compounds of the formula (I) may be prepared by reacting a compound of the formula



wherein Ar, R^1 and R^2 are defined as above and Z represents a halogen atom or an organic sulfonyloxy group (e.g. tosyloxy, mesyloxy, etc.), with a compound of the formula



wherein Am is defined as above.

The reaction may be carried out usually in a solvent such as methanol, ethanol, isopropanol, benzene, toluene, xylene, dimethylformamide, chloroform, dichloroethane, acetone, methyl ethyl ketone, etc., at a temperature between room temperature and 140°C , preferably

between 50°C and 110°C, in the presence of potassium carbonate, sodium carbonate, triethylamine or like acid acceptor, for 1 to 48 hours, preferably 4 to 18 hours. The reaction may be accelerated by the use of a catalyst.

- 5 Examples of such catalyst are potassium iodide, sodium iodide, etc.

The compound of the formula (I) may be converted into an acid addition salt. Typical examples of such an acid addition salt which is pharmaceutically acceptable
10 are salts formed with use of hydrochloric acid, sulfuric acid, phosphoric acid, methanesulfonic acid, maleic acid, oxalic acid, succinic acid, fumaric acid, acetic acid, lactic acid and citric acid.

The experiments carried out for demonstrating
15 anti-apomorphine activity of the compounds of this invention in mice will be described below.

Experimental method:

Groups of 5 male dd-mice (20-25 g body weight) each were used. Apomorphine hydrochloride (0.5 mg/kg)
20 was subcutaneously administered 60 minutes after oral administration of test compound. Immediately after the apomorphine treatment, motor activity was determined for 20 minutes by animex. For the control groups, 0.5% methylcellulose solution was administered instead of
25 test compound. The ED₅₀, a dose which inhibited the


motor activity by 50% as compared with the control, was determined.

Results:


Compound	Anti-apomorphine activity ED ₅₀ (mg/kg, p.o.)
5 A	1.7
B	2.1
C	3.4
Clozapine	10

- 10 A: 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-oxo-1-benzimidazoliny)l)piperidine fumarate
- B: 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-fluoro-2-oxo-1-benzimidazoliny)l)-piperidine maleate
- 15 C: 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorophenyl)-4-hydroxypiperidine fumarate

20 The compounds of the formula (II) are novel and may be prepared, for example, by reducing a compound of the formula



wherein Ar, R¹ and R² are as defined above and R represents a lower alkyl group, with use of sodium borohydride etc., and reacting the resulting compound of the formula



5 wherein Ar, R¹ and R² are as defined above, with thionyl chloride, phosphorus tribromide or like halogenating agent or tosyl chloride, mesyl chloride or like organic sulfonating agent.

Reference Example 1

10 To a solution of 7 g of ethyl 5-phenyl-4,5-dihydroisoxazol-3-ylcarboxylate in 70 ml of methanol, cooled with ice, is added 1.5 g of sodium borohydride in small portions with stirring. After 4 hours has passed, the solvent is distilled off under reduced pressure and the residue is extracted with ethyl acetate. The extract
15 is washed with water, dried and the solvent is evaporated. The crustalline residue thus obtained is recrystallized from isopropyl ether, giving 3-hydroxymethyl-5-phenyl-4,5-dihydroisoxazole in the form of white crystals. Melting point : 73-74°C.

20

Reference Example 2

3-Hydroxymethyl-5-phenyl-4,5-dihydroisoxazole
(3.2 g) is dissolved in 50 ml of anhydrous ether. To
the solution cooled with ice is slowly added dropwise
2.6 g of thionyl chloride with stirring. The reaction
5 mixture is allowed to stand overnight at room tem-
perature and then the solvent is distilled off to give
3-chloromethyl-5-phenyl-4,5-dihydroisoxazole in the form
of yellow brown oil.

The compounds of the formula (I) are used in
10 combination with a suitable and conventional pharma-
ceutically acceptable excipient in the form of a pharma-
ceutical composition. The pharmaceutical composition
may take usual forms such as of tablets, capsules,
powders, granules, injection solutions, etc.

15 When administered for pharmaceutical uses, the
compounds of this invention may, for example, be formulated
into a pharmaceutical composition as follows.

Tablets (10 mg) may be prepared from the
following ingredients:

20	Compound (I) or salt thereof	10 mg
	Lactose	53 mg
	Crystalline cellulose	15 mg
	Corn starch	20 mg
	Polyvinyl alcohol	1.5 mg
25	Magnesium stearate	0.5 mg

100 mg

A compound (I) or salt thereof, crystalline cellulose and corn starch are mixed together and then the mixture is kneaded with 5% polyvinyl alcohol. The resulting mixture is granulated, dried and the dry granules are
5 passed through 24-mesh screen. The fine granules are mixed with magnesium stearate to form granules for the preparation of tablets. Tablets are prepared by compressing the granules on punches (6.5 mm, 7.0R).

The dose of the compounds of the formula (I)
10 ranges from 0.005 to 100 mg/kg body weight/day, preferably from 0.01 to 50 mg/kg body weight/day, which may be administered at one time or at several times, although variable depending on the age, body weight and/or severity of the conditions to be treated or response to the
15 medication.

This invention will be better understood from the following examples, which are not to be construed as limitative of the present invention.

Example 1

20 3-Chloromethyl-5-phenyl-4,5-dihydroisoxazole (5.87 g), 6.3 g of 4-(4-chlorophenyl)-4-hydroxypiperidine, 4 g of potassium carbonate and 50 ml of ethanol are heated to 60-70°C with stirring for 6 hours. The reaction mixture is filtered and the filtrate is condensed by
25 distillation under reduced pressure. To the residue are

added 200 ml of ethyl acetate and 100 ml of water. The organic layer is separated off, washed with water, dried on magnesium sulfate and evaporated under reduced pressure. The residue thus obtained is dissolved in

5 isopropyl ether and alcoholic hydrochloric acid is added to the solution. The crystals thus formed is filtered and then recrystallized from isopropyl alcohol, giving 1-(5-phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(4-chlorophenyl)-4-hydroxypiperidine hydrochloride.

10 Melting point: 175-176°C (decomposition)

Example 2

3-Chloromethyl-5-(4-fluorophenyl)-4,5-dihydroisoxazole (40 g), 52 g of 4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine, 30 g of potassium carbonate,

15 15 g of potassium iodide and 1 liter of ethanol are heated to a temperature of 70 to 75°C with stirring for 48 hours. The reaction mixture is then filtered and the mother liquor is condensed under reduced pressure. To the residue are added 800 ml of chloroform and 500 ml of

20 water and the mixture is stirred. The organic layer is separated off, washed with water and dried on magnesium sulfate, and the solvent is distilled off. To the resulting residue are added 130 ml of acetone and 100 ml of isopropyl ether. The crystals thus precipitated are

25 filtered and recrystallized from a mixture of acetone

(400 ml) and isopropyl ether (450 ml), to give 67.5 g of 1-[5-(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine having a melting point of 163 to 164°C. Hydrochloride of this
5 compound melts at 216°C (decomposition).

A 46.5 g-quantity of the above compound (free base) is dissolved in 200 ml of ethanol, and a solution of 15 g of L-tartaric acid in 200 ml of water is added to the ethanol solution. The resulting mixture is
10 allowed to stand at room temperature. The crystals thus precipitated is recrystallized three times from ethanol-water (6:4) to give tartrate monohydrate as colorless prisms. The tartrate monohydrate is treated with an aqueous solution of sodium bicarbonate to give (-)-1-[5-
15 (4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine. Melting point: 143-145°C. $[\alpha]_D^{25}$: - 108.8 (chloroform).

The (+)-isomer of the above compound is obtained in the same manner as above with use of D-tartaric acid.
20 Melting point: 142-144°C. $[\alpha]_D^{25}$: + 112.6 (chloroform).

Example 3

3-Chloromethyl-5-(4-fluorophenyl-4,5-dihydroisoxazole (3.2 g), 3.5 g of 4-oxo-1-phenyl-1,3,8-
25 triazaspiro[4,5]decane, 2.1 g of potassium carbonate and

100 ml of ethanol are refluxed for 7.5 hours with stirring. The resulting reaction mixture is filtered and the mother liquor is concentrated. To the residue obtained is added 100 ml of water and the mixture is extracted
5 with ethyl acetate. The extract is washed with water and dried on magnesium sulfate and the solvent is distilled off under reduced pressure. The resulting residue is dissolved in a small amount of alcohol and alcoholic hydrochloric acid is added to the solution. The crystals
10 thus precipitated are filtered and recrystallized from methanol to give 5-(4-fluorophenyl)-3-(4-oxo-1-phenyl-1,3,8-triazaspiro[4,5]decan-8-ylmethyl)-4,5-dihydroisoxazole hydrochloride. Melting point: 219°C (decomposition).

The following compounds may be prepared in the
15 same manner as in the preceding Examples.

- ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorophenyl)-4-hydroxypiperidine

Melting point of 1/2 fumarate: 147-148°C

- ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
20 4-(2-oxo-1-benzimidazoliny1)piperidine

Melting point of fumarate: 206°C (decomp.)

- ° 1-[5-(2-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-chloro-2-oxo-1-benzimidazoliny1)piperidine

Melting point of hydrochloride: 244°C (decomp.)

25

- ° 1-[5-(3-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-chloro-2-oxo-1-benzimidazoliny1)piperidine
Melting point of maleate: 197°C (decomp.)
- 5 ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-fluoro-2-oxo-1-benzimidazoliny1)piperidine
Melting point of maleate: 201°C (decomp.)
- ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-carbamoyl-4-piperidino-piperidine
Melting point of dihydrochloride: 234°C (decomp.)
- 10 ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-benzylpiperidine
Melting point of hydrochloride: 184°C
- ° 1-(5-Methyl-5-phenyl-4,5-dihydroisoxazol-3-ylmethyl)-
4-(5-chloro-2-oxo-1-benzimidazoliny1)piperidine
15 Melting point of maleate: 218°C (decomp.)
- ° 5-Phenyl-3-[4-oxo-1-(4-bromophenyl)-1,3,8-triazaspiro-
[4,5]-decan-8-ylmethyl]-4,5-dihydroisoxazole
Melting point of maleate: 221°C (decomp.)
- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-
20 hydroxypiperidine
Melting point of maleate: 115-119°C
- ° 1'-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-
1,2,3,5,6,7,8,8a-octahydro-2-oxoimidazo[1,2-a]-
pyridine-3-spiro-4'-piperidine
25 Melting point of dihydrochloride: 223°C (decomp.)

- ° 1-[5-(4-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-chloro-2-oxo-1-benzimidazoliny1)piperidine
Melting point of hydrochloride: 230°C (decomp.)
- 5 ° 1-[5-(4-Methoxyphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-chloro-2-oxo-1-benzimidazoliny1)piperidine
Melting point of hydrochloride: 229°C (decomp.)
- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-
chloro-2-thioxo-1-benzimidazoliny1)piperidine
Melting point of 1/2 fumarate: 208-209°C
- 10 ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-
methoxypiperidine
Melting point of hydrochloride: 162-164°C
- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-
carbamoyl-4-piperidino-piperidine
15 Melting point of dihydrochloride: 158°C (decomp.)
- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-
benzylpiperidine
Melting point of hydrochloride: 208°C (decomp.)
- 20 ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(2-oxo-
1-benzimidazoliny1)piperidine
Melting point of fumarate: 206°C (decomp.)
- ° 1-[5-(2-Pyridyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-
(5-chloro-2-oxo-1-benzimidazoliny1)piperidine
Melting point of maleate: 188°C

- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-fluoro-2-oxo-1-benzimidazoliny)l)piperidine
Melting point of hydrochloride: 229°C (decomp.)
- 5 ° 5-(Phenyl-3-(4-oxo-1-phenyl-1,3,8-triazaspiro[4,5]-decan-8-ylmethyl)-4,5-dihydroisoxazole
Melting point of hydrochloride: 226°C (decomp.)
- ° 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine
Melting point of maleate: 204°C (decomp.)
- 10 ° 1-(5-Phenyl-3-isoxazoliny)methyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine
Melting point of hydrochloride: 250°C (decomp.)
- ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-imidazole
15 Melting point of fumarate: 110-111°C
- ° 1-[5-(2-Pyridyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(3-methyl-2,4-dioxo-1-hexahydropyrimidinyl)piperidine
Melting point of maleate: 169°C (decomp.)
- ° 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine
20
- ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-thioxo-1-benzimidazoliny)l)piperidine
Melting point of fumarate: 125°C (decomp.)
- ° 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(2-thioxo-1-benzimidazoliny)l)piperidine
25

- ° 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-
4-(2-oxo-1-benzimidazoliny1)piperidine

Melting point: 197-199°C

- 5 ° 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-
4-(5-fluoro-2-oxo-1-benzimidazoliny1)piperidine

- ° 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-
4-hydroxy-4-(4-chlorophenyl)piperidine

Melting point: 153-154°C

- 10 ° 1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny1)-
piperidine

Melting point: 199-200°C

- ° 1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-(2-thioxo-1-benzimidazoliny1)piperidine

- 15 ° 1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-(2-oxo-1-benzimidazoliny1)piperidine

- ° 1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-(5-fluoro-2-oxo-1-benzimidazoliny1)-
piperidine

- 20 ° 1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-hydroxy-4-(4-chlorophenyl)piperidine

Melting point: 163-164°C

- 25 ° 1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-
ylmethyl]-4-(4-chloro-3-trifluoromethylphenyl)-
4-hydroxypiperidine

° 3-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-2,3,4,4a,5,6-hexahydro-1H-pyrazino-[1,2-a]-quinoline


Melting point of oxalate: 148°C (decomp.)

5










While the invention has been described in detail and with reference to specific embodiments thereof, it is apparent that various alterations and modifications can be made without departing from the spirit and scope thereof.

CLAIMS:


1. An isoxazole derivative represented by the formula



or salts thereof wherein Ar represents a phenyl group
5 which may optionally be substituted with a halogen atom
or a lower alkoxy group, or a pyridyl group R¹ represents
a hydrogen atom, a lower alkyl or a group represented by
a group Ar, R² represents a hydrogen atom, or alternatively
R¹ and R² are bound together and form a carbon-carbon
10 bond, and Am represents an amino residue selected from
the group consisting of the following residues:



and



wherein R^3 represents a hydrogen atom or a lower alkyl group, X^1 and X^2 each represent a hydrogen atom, a halogen atom or a trifluoromethyl group and Y represents O or S.

5

2. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-chloro-2-oxo-1-benzimidazoliny)l)piperidine.

3. The compound of claim 1:

10

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(2-oxo-1-benzimidazoliny)l)piperidine.

4. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(5-fluoro-2-oxo-1-benzimidazoliny)l)piperidine.

5. The compound of claim 1:

15

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(4-chlorophenyl)-4-hydroxypiperidine.

6. The compound of claim 1:

5-(4-Fluorophenyl)-3-(4-oxo-1-phenyl-1,3,8-triazaspiro-
[4,5]-decan-8-ylmethyl)-4,5-dihydroisoxazole.

20

7. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-
4-(2-thioxo-1-benzimidazoliny)l)piperidine.

8. The compound of claim 1:

25








1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl-4-(5-chloro-
2-thioxo-1-benzimidazoliny)l)piperidine.

0040639


- 20 -

9. A pharmaceutical composition comprising
a compound of claim 1 and a pharmaceutically acceptable
excipient.

21



and



These compounds are useful as drugs such as psychotropic agent and antiemetic agent.

International Application No

1. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ^a

According to International Patent Classification (IPC) or to both National Classification and IPC

II. FIELDS SEARCHED

Minimum Documentation Searched 4

Classification System

Classification Symbols

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched 5

III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴

Category *	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
------------	--	-------------------------------------

* Special categories of cited documents: 18

"A" document defining the general state of the art

"E" earlier document but published on or after the international filing date

"L" document cited for special reason other than those referred to in the other categories

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but on or after the priority date claimed

"T" later document published on or after the international filing date or priority date and not in conflict with the application, but cited to understand the principle or theory underlying the invention

"X" document of particular relevance

IV. CERTIFICATION

Date of the Actual Completion of the International Search :

Date of Mailing of this International Search Report :

February 20, 1981 (20.02.81)

March 2, 1981 (02.03.81)

International Searching Authority 1

Signature of Authorized Officer 20

Japanese Patent Office

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 040 639
B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 30.05.84

(21) Application number: 80902298.1

(22) Date of filing: 25.11.80

(38) International application number:
PCT/JP80/00286

(17) International publication number:
WO 81/01554 11.06.81 Gazette 81/14

(51) Int. Cl.³: **C 07 D 413/06,**
C 07 D 413/14,
C 07 D 471/04,
C 07 D 471/10, C 07 D 471/20

(54) **ISOXAZOLE DERIVATIVES.**

(30) Priority: 28.11.79 JP 154715/79

(43) Date of publication of application:
02.12.81 Bulletin 81/48

(45) Publication of the grant of the patent:
30.05.84 Bulletin 84/22

(84) Designated Contracting States:
DE FR GB NL

(50) References cited:
DE-A-2 245 971
JP-A-49 024 973
US-A-3 890 323
US-A-4 133 889

(70) Proprietor: Yoshitomi Pharmaceutical
Industries, Ltd.
35 Hiranomachi 3-chome Higashi-ku
Osaka-shi Osaka 541 (JP)

(72) Inventor: KAWAKITA, Takeshi
7-8, Chuoma-chi 1-chome
Nakatsu-shi, Oita 871 (JP)
Inventor: MURO, Tomio
631-7, Oaza-kakize Nakatsu-shi
Oita 871 (JP)
Inventor: SETOYUCHI, Michihide
5-21, Chuomachi 2-chome
Nakatsu-shi, Oita 871 (JP)

(74) Representative: Heunemann, Dieter Dr.
VOSSIUS VOSSIUS TAUCHNER HEUNEMANN
RAUH P.O. Box 86 07 67 Siebertstrasse 4
D-8000 Munich 86 (DE)

EP 0 040 639 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).

Courier Press, Leamington Spa, England.

Description










Technical Field and Disclosure of the Invention

This invention relates to novel isoxazole derivatives having spontaneous locomotor suppressing activity, anti-apomorphine activity and similar activity which are useful as drugs such as psychotropic and antiemetic agents and are represented by the general formula




15 to pharmaceutically acceptable acid addition salts thereof, and to a process for preparing these compounds.

In the foregoing formula, Ar represents a phenyl group optionally containing a lower alkoxy group or a halogen atom as a substituent, or a pyridyl group, R¹ represents a hydrogen atom, a lower alkyl group or a group represented by Ar, R² represents a hydrogen atom or alternatively R¹ and R² are bound together and form a carbon-carbon bond, and Am represents an amino residue selected from the group consisting of the following residues:



and




wherein R³ represents a hydrogen atom or a lower alkyl group, X¹ and X² each represent a hydrogen atom, a halogen atom or a trifluoromethyl group, and Y represents O or S.

The term "halogen" herein includes fluorine, chlorine and bromine. The term "lower alkoxy" herein represents a methoxy, ethoxy, propoxy and butoxy. The term "lower alkyl" herein represents methyl, ethyl, propyl and butyl.

DE—OS 2 245 971 discloses isoxazole derivatives that are intermediates for preparing pharmaceutical substances. US—PS 3 890 323 discloses phenylketone derivatives that show central nervous system depressing activity.

The compounds of the formula (I) may be prepared by reacting a compound of the formula



wherein Ar, R¹ and R² are as defined above and Z represents a halogen atom or an organic sulfonyloxy group (e.g. tosyloxy or mesyloxy), with a compound of the formula



wherein Am is as defined above.

The reaction may be carried out usually in a solvent such as methanol, ethanol, isopropanol, benzene, toluene, xylene, dimethylformamide, chloroform, dichloroethane, acetone or methyl ethyl ketone, at a temperature between room temperature and 140°C, preferably between 50°C and 110°C, in the presence of potassium carbonate, sodium carbonate, triethylamine or other acid acceptors, for 1 to 48 hours, preferably 4 to 18 hours. The reaction may be accelerated by the use of a catalyst. Examples of such catalyst are potassium iodide and sodium iodide.

The compound of the formula (I) may be converted into an acid addition salt. Typical examples of such an acid addition salt which is pharmaceutically acceptable are salts formed with the aid of hydrochloric acid, sulfuric acid, phosphoric acid, methanesulfonic acid, maleic acid, oxalic acid, succinic acid, fumaric acid, acetic acid, lactic acid and citric acid.

The experiments carried out for demonstrating the anti-apomorphine activity of the compounds of this invention in mice will be described below.

Experimental method:

Groups of 5 male dd-mice (20—25 g body weight) each were used. Apomorphine hydrochloride (0.5 mg/kg) was subcutaneously administered 60 minutes after oral administration of the test compound. Immediately after the apomorphine treatment, motor activity was determined for 20 minutes by animex. For the control groups, 0.5% methylcellulose solution was administered instead of the test compound. The ED₅₀, a dose which inhibited the motor activity by 50% was compared with the control, was determined.

Results:

Compound	Anti-apomorphine activity ED ₅₀ (mg/kg, p.o.)
A	1.7
B	2.1
C	3.4
Clozapine	10


A: 1 - [5 - (4 - Fluorophenyl) - 4,5 - dihydroisoxazol - 3 - ylmethyl] - 4 - (2 - oxo - 1 - benzimidazoliny)l)piperidine fumarate

B: 1 - [5 - (4 - Fluorophenyl) - 4,5 - dihydroisoxazol - 3 - ylmethyl] - 4 - (5 - fluoro - 2 - oxo - 1 - benzimidazoliny)l)piperidine maleate


C: 1 - [5 - (4 - Fluorophenyl) - 4,5 - dihydroisoxazol - 3 - ylmethyl] - 4 - (4 - chlorophenyl) - 4 - hydroxypiperidine fumarate

0 040 639

The compounds of the formula (II) are novel and may be prepared, for example, by reducing a compound of the formula



wherein Ar, R¹ and R² are as defined above and R represents a lower alkyl group, using sodium borohydride and reacting the resulting compound of the formula



wherein Ar, R¹ and R² are as defined above, with thionyl chloride, phosphorus tribromide or similar halogenating agents or tosyl chloride, mesyl chloride or similar organic sulfonating agents.

Reference Example 1

To a solution of 7 g of ethyl 5-phenyl-4,5-dihydroisoxazol-3-ylcarboxylate in 70 ml of methanol, cooled with ice, are added 1.5 g of sodium borohydride in small portions with stirring. After 4 hours have passed, the solvent is distilled off under reduced pressure and the residue is extracted with ethyl acetate. The extract is washed with water, dried and the solvent is evaporated. The crystalline residue thus obtained is recrystallized from isopropyl ether, giving 3-hydroxymethyl-5-phenyl-4,5-dihydroisoxazole in the form of white crystals. Melting point: 73—74°C.

Reference Example 2

3-Hydroxymethyl-5-phenyl-4,5-dihydroisoxazole (3.2 g) is dissolved in 50 ml of anhydrous ether. To the solution cooled with ice are added slowly, dropwise 2.6 g of thionyl chloride under stirring. The reaction mixture is allowed to stand overnight at room temperature and the solvent is then distilled off to give 3-chloromethyl-5-phenyl-4,5-dihydroisoxazole in the form of a yellow brown oil.

The compounds of the formula (I) are used in combination with a suitable and conventional pharmaceutically acceptable excipient in the form of a pharmaceutical composition. The pharmaceutical composition may be configured in the usual form such as tablets, capsules, powders, granules or injection solutions.

When administered for pharmaceutical uses, the compounds of this invention may, for example, be formulated into a pharmaceutical composition as follows.

Tablets (10 mg) may be prepared from the following ingredients:

Compound (I) or salt thereof	10 mg
Lactose	53 mg
Crystalline cellulose	15 mg
Corn starch	20 mg
Polyvinyl alcohol	1.5 mg
Magnesium stearate	0.5 mg
	100 mg

A compound (I) or a salt thereof, crystalline cellulose and corn starch are mixed together and the mixture is then kneaded with 5% polyvinyl alcohol. The resulting mixture is granulated, dried and the dry granules are passed through a 24-mesh screen. The fine granules are mixed with magnesium stearate to form granules for the preparation of tablets. Tablets are prepared by compressing the granules on punches (6.5 mm, 7.0R).

The dose of the compounds of the formula (I) ranges from 0.005 to 100 mg/kg body weight/day, preferably from 0.01 to 50 mg/kg body weight/day, which may be administered at one time or at

several times, although variable depending on the age, body weight and/or severity of the conditions to be treated or response to the medication.

This invention will be better understood from the following examples.

Example 1

3 - Chloromethyl - 5 - phenyl - 4,5 - dihydroisoxazole (5.87 g), 6.3 g of 4 - (4 - chlorophenyl) - 4 - hydroxypiperidine, 4 g of potassium carbonate and 50 ml of ethanol are heated to 60—70°C under stirring for 6 hours. The reaction mixture is filtered and the filtrate is condensed by distillation under reduced pressure. To the residue are added 200 ml of ethyl acetate and 100 ml of water. The organic layer is separated off, washed with water, dried on magnesium sulfate and evaporated under reduced pressure. The residue thus obtained is dissolved in isopropyl ether and alcoholic hydrochloric acid is added to the solution. The crystals thus formed are filtered and then recrystallized from isopropyl alcohol, giving 1-[5-phenyl-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorophenyl)-4-hydroxypiperidine hydrochloride. Melting point: 175—176° (decomposition).

Example 2

3 - Chloromethyl - 5 - (4 - fluorophenyl) - 4,5 - dihydroisoxazole (40 g), 52 g of 4 - (5 - chloro-2 - oxo - 1 - benzimidazolyl)piperidine, 30 g of potassium carbonate, 15 g of potassium iodide and 1 liter of ethanol are heated to a temperature of 70 to 75°C with stirring for 48 hours. The reaction mixture is then filtered and the mother liquor is condensed under reduced pressure. To the residue are added 800 ml of chloroform and 500 ml of water and the mixture is stirred. The organic layer is separated off, washed with water and dried on magnesium sulfate, and the solvent is distilled off. To the resulting residue are added 130 ml of acetone and 100 ml of isopropyl ether. The crystals thus precipitated are filtered and recrystallized from a mixture of acetone (400 ml) and isopropyl ether (450 ml), to give 67.5 g of 1-[5-(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazolyl)piperidine having a melting point of 163 to 164°C. The hydrochloride of this compound melts at 216°C (decomposition).

A 46.5 g-quantity of the above compound (free base) is dissolved in 200 ml of ethanol, and a solution of 15 g of L-tartaric acid in 200 ml of water is added to the ethanol solution. The resulting mixture is allowed to stand at room temperature. The crystals thus precipitated are recrystallized three times from ethanol-water (6:4) to give tartrate monohydrate as colorless prisms. The tartrate monohydrate is treated with an aqueous solution of sodium bicarbonate to give (-)-1-[5-(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazolyl)piperidine. Melting point: 143—145°C. $[\alpha]_D^{25}$: - 108.8 (chloroform).

The (+)-isomer of the above compound is obtained in the same manner as above using D-tartaric acid. Melting point: 142—144°C. $[\alpha]_D^{25}$: +112.6 (chloroform).

Example 3

3 - Chloromethyl - 5 - (4 - fluorophenyl) - 4,5 - dihydroisoxazole (3.2 g), 3.5 g of 4 - oxo - 1 - phenyl-1,3,8-triazaspiro[4,5]decan-2-one, 2.1 g of potassium carbonate and 100 ml of ethanol are refluxed for 7.5 hours while stirring. The resulting reaction mixture is filtered and the mother liquor is concentrated. To the residue obtained are added 100 ml of water and the mixture is extracted with ethyl acetate. The extract is washed with water and dried on magnesium sulfate and the solvent is distilled off under reduced pressure. The resulting residue is dissolved in a small amount of alcohol and alcoholic hydrochloric acid is added to the solution. The crystals thus precipitated are filtered and recrystallized from methanol to give 5-(4-fluorophenyl)-3-(4-oxo-1-phenyl-1,3,8-triazaspiro[4,5]decan-8-ylmethyl)-4,5-dihydroisoxazole hydrochloride. Melting point: 219°C (decomposition).

The following compounds may be prepared in the same manner as in the preceding Examples.

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorophenyl)-4-hydroxypiperidine
Melting point of 1/2 fumarate: 147—148°C

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-oxo-1-benzimidazolyl)piperidine
Melting point of fumarate: 206°C (decomp.)

1-[5-(4-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazolyl)-piperidine

Melting point of hydrochloride: 244°C (decomp.)

1-[5-(3-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazolyl)-piperidine

Melting point of maleate: 197°C (decomp.)

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-fluoro-2-oxo-1-benzimidazolyl)-piperidine

Melting point of maleate: 201°C (decomp.)

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-carbamoyl-piperidino-piperidine

Melting point of dihydrochloride: 234°C (decomp.)

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-benzylpiperidine

Melting point of hydrochloride: 184°C

- 1-(5-Methyl-5-phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)-piperidine
Melting point of maleate: 218°C (decomp.)
5-Phenyl-3-[4-oxo-1-(4-bromophenyl)-1,3,8-triazaspiro-[4,5]-decan-8-ylmethyl]-4,5-dihydroisoxazole
5 isoxazole
Melting point of maleate: 221°C (decomp.)
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-hydroxypiperidine
Melting point of maleate: 115—119°C
1'-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-1,2,3,5,6,7,8,8a-octahydro-2-oxoimidazo[1,2-a]-pyridine-3-spiro-4'-piperidine
10 pyridine-3-spiro-4'-piperidine
Melting point of dihydrochloride: 223°C (decomp.)
1-[5-(4-Chlorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)-piperidine
Melting point of hydrochloride: 230°C (decomp.)
15 1-[5-(4-Methoxyphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)-piperidine
Melting point of hydrochloride: 229°C (decomp.)
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-thioxo-1-benzimidazoliny)piperidine
Melting point of 1/2 fumarate: 208—209°C
20 1-(5-Phenyl-4,5-dihydroisoxazole-3-ylmethyl)-4-methoxypiperidine
Melting point of hydrochloride: 162—164°C
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-carbamoyl-4-piperidino-piperidine
Melting point of dihydrochloride: 158°C (decomp.)
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-benzylpiperidine
25 Melting point of hydrochloride: 208°C (decomp.)
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(2-oxo-1-benzimidazoliny)piperidine
Melting point of fumarate: 206°C (decomp.)
1-[5-(2-Pyridyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)piperidine
idine
30 Melting point of maleate: 188°C
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-fluoro-2-oxo-1-benzimidazoliny)piperidine
Melting point of hydrochloride: 229°C (decomp.)
5-(Phenyl-3-(4-oxo-1-phenyl-1,3,8-triazaspiro[4,5]-decan-8-ylmethyl)-4,5-dihydroisoxazole
Melting point of hydrochloride: 226°C (decomp.)
35 1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)piperidine
Melting point of maleate: 204°C (decomp.)
1-(5-Phenyl-3-isoxazolinylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)piperidine
Melting point of hydrochloride: 250°C (decomp.)
1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-imidazole
40 Melting point of fumarate: 110—111°C
1-[5-(2-Pyridyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(3-methyl-2,4-dioxo-1-hexahydropyrimidinyl)piperidine
Melting point of maleate: 169°C (decomp.)
1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chloro-2-oxo-1-benzimidazoliny)-piperidine
45 piperidine
1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-thioxo-1-benzimidazoliny)-piperidine
Melting point of fumarate: 125°C (decomp.)
1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(2-thioxo-1-benzimidazoliny)piperidine
50 1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(2-oxo-1-benzimidazoliny)piperidine
Melting point: 197—199°C
1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-fluoro-2-oxo-1-benzimidazoliny)-piperidine
1-(5,5-Diphenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-hydroxy-4-(4-chlorophenyl)piperidine
55 Melting point: 153—154°C
1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)piperidine
Melting point: 199—200°C
1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-thioxo-1-benzimidazoliny)-piperidine
60 piperidine
1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-oxo-1-benzimidazoliny)piperidine
idine
1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-fluoro-2-oxo-1-benzimidazoliny)piperidine
65

1-[5,5-Bis(4-fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-hydroxy-4-(4-chlorophenyl)-piperidine

Melting point: 163—164°C


1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chloro-3-trifluoromethylphenyl)-4-hydroxypiperidine

3-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-2,3,4,4a,5,6-hexahydro-1H-pyrazino-[1,2-a]-quinoline

Melting point of oxalate: 148°C (decomp.)


10 Claims

1. An isoxazole derivative represented by the formula



20 or salts thereof wherein Ar represents a phenyl group which may optionally be substituted with a halogen atom or a C₁₋₄ alkoxy group, or a pyridyl group R¹ represents a hydrogen atom, a C₁₋₄ alkyl or a group represented by a group Ar, R² represents a hydrogen atom, or alternatively R¹ and R² are bound together and form a carbon-carbon bond, and Am represents an amino residue selected from the group

25 consisting of the following residues:



wherein R^3 represents a hydrogen atom or a C_{1-4} alkyl group, X^1 and X^2 each represent a hydrogen atom, a halogen atom or a trifluoromethyl group and Y represents O or S.

2. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-oxo-1-benzimidazolyl)-piperidine.

3. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-oxo-1-benzimidazolyl)piperidine.

4. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-fluoro-2-oxo-1-benzimidazolyl)-piperidine.

5. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorophenyl)-4-hydroxypiperidine.

6. The compound of claim 1:

5-(4-Fluorophenyl)-3-(4-oxo-1-phenyl-1,3,8-triazaspiro-[4,5]-decan-8-ylmethyl)-4,5-dihydroisoxazole.

7. The compound of claim 1:

1-[5-(4-Fluorophenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-thioxo-1-benzimidazolyl)-piperidine.


8. The compound of claim 1:

1-[5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chloro-2-thioxo-1-benzimidazolyl)piperidine.


9. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable excipient.

Revendications

1. Dérivé isoxazole, caractérisé en ce qu'il est représenté par la formule générale ci-après



ou ses sels, dans laquelle Ar représente un groupe phényle qui peut éventuellement être substitué avec un atome d'halogène ou un groupe alkoxy en C_1 à 4 , ou un groupe pyridyle, R^1 représente un atome d'hydrogène, un radical alkyle en C_1 à 4 ou un groupe représenté par le groupe Ar, R^2 représente un atome d'hydrogène, ou, dans une variante, R^1 et R^2 peuvent être reliés et former une liaison carbone-carbone, et Am représente un résidu amino choisi dans le groupe formé par les résidus suivants:




où R³ représente un atome d'hydrogène ou un groupe alkyle en C₁ à ₄, X¹ et X² sont chacun un atome d'hydrogène, un atome d'halogène ou un groupe trifluorométhyle et Y représente un atome de —O— ou —S—.

2. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-(4-fluorophényl)-4,5-dihydroisoxazol-3-ylméthyl]-4-(5-chloro-2-oxo-1-benzimidazoliny)-
pipéridine.
3. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-(4-fluorophényl)-4,5-dihydroisoxazol-3-ylméthyl]-4-(2-oxo-1-benzimidazoliny)pipéridine.
4. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-(4-fluorophényl)-4,5-dihydroisoxazol-3-ylméthyl]-4-(5-fluoro-2-oxo-1-benzimidazoliny)-
pipéridine.
5. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-(4-fluorophényl)-4,5-dihydroisoxazol-3-ylméthyl]-4-(4-chlorophényl)-4-hydroxypipéridine.
6. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
5-(4-fluorophényl)-3-(4-oxo-1-phényl-1,3,8-triazaspiro-[4,5]-décan-8-ylméthyl)-4,5-dihydroiso-
xazole.
7. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-(4-fluorophényl)-4,5-dihydroisoxazol-3-ylméthyl]-4-(2-thio-1-benzimidazoliny)-
pipéridine.
8. Composé selon la revendication 1, caractérisé en ce qu'il s'agit de la:
1-[5-phényl-4,5-dihydroisoxazol-3-ylméthyl]-4-(5-chloro-2-thio-1-benzimidazoliny)pipéridine.
9. Composition pharmaceutique, caractérisée en ce qu'elle comprend un composé selon la
revendication 1 et un excipient acceptable du point de vue pharmaceutique.

Patentansprüche

1. Ein Isoxazolderivat der Formel

5




(I)

10


oder seine Salze, in der Ar eine gegebenenfalls durch ein Halogenatom oder einen C₁₋₄-Alkoxygruppe substituierte Phenylgruppe oder eine Pyridylgruppe bedeutet, R¹ ein Wasserstoffatom, eine C₁₋₄-Alkylgruppe oder den Rest Ar bedeutet, R² ein Wasserstoffatom darstellt, oder R¹ und R² zusammen eine Kohlenstoff-Kohlenstoffbindung darstellen, und Am einen Aminorest bedeutet, der ausgewählt ist aus der Gruppe der folgenden Reste:

15



20




25




30




35




40




45




50



55



und



60

in denen R³ ein Wasserstoffatom oder eine C₁₋₄-Alkylgruppe bedeutet, X¹ und X² jeweils ein Wasserstoffatom, ein Halogenatom oder eine Trifluormethylgruppe darstellen und Y O oder S bedeutet.

2. Die Verbindung nach Anspruch 1:

1-[5-(4-Fluorphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-chlor-2-oxo-1-benzimidazolyl)-piperidin.

65

3. Die Verbindung nach Anspruch 1:
1-[5-(4-Fluorphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-oxo-1-benzimidazoliny)piperidin.
4. Die Verbindung nach Anspruch 1:
1-[5-(4-Fluorphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(5-fluor-2-oxo-1-benzimidazoliny)-
5 piperidin.
5. Die Verbindung nach Anspruch 1:
1-[5-(4-Fluorphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(4-chlorphenyl)-4-hydroxypiperidin.
6. Die Verbindung nach Anspruch 1:
5-(4-Fluorphenyl)-3-(4-oxo-1-phenyl-1,3,8-triazaspiro-[4,5]-decan-8-ylmethyl)-4,5-dihydroiso-
10 xazol.
7. Die Verbindung nach Anspruch 1:
1-[5-(4-Fluorphenyl)-4,5-dihydroisoxazol-3-ylmethyl]-4-(2-thioxo-1-benzimidazoliny)piperidin.
8. Die Verbindung nach Anspruch 1:
1-(5-Phenyl-4,5-dihydroisoxazol-3-ylmethyl)-4-(5-chlor-2-thioxo-1-benzimidazoliny)piperidin.
- 15 9. Ein Arzneimittel, enthaltend eine Verbindung nach Anspruch 1 und einen pharmazeutisch ver-
träglichen Trägerstoff.

20

25

30

35

40

45

50

55

60

65

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.